

6

Vibrational Stark Effects of Nitriles

I. Methods and Experimental Results

Reprinted with permission from:

Andrews, S.S.; Boxer, S.G. *J. Phys. Chem. A* **2000**, *104*, 11853.

Copyright 2000 American Chemical Society.

In many ways the most valuable application of the data of infrared and Raman studies is to the calculation of the heat capacity, entropy, and free energy of gaseous molecules.

— Wilson, Decius, and Cross

*Molecular Vibrations. The Theory of
Infrared and Raman Vibrational Spectra*

Abstract

Vibrational Stark effects were measured for the nitrile stretch mode of several nitriles in frozen 2-methyl-tetrahydrofuran glass. Samples included unconjugated aliphatic nitriles, conjugated and unconjugated aliphatic dinitriles, and aromatic mono- and dinitriles. Difference dipole moments, $\Delta\mu$, equivalent to the linear Stark tuning rate, range from $0.01/f$ to $0.04/f$ Debye ($0.2/f$ to $0.7/f$ $\text{cm}^{-1}/(\text{MV}/\text{cm})$) for most samples studied, with aromatic compounds towards the high end and symmetric dinitriles towards the low end (the local field correction factor, f , is expected to be similar for all samples in the same solvent). Transition dipoles and $\Delta\mu$ values for aromatic nitriles correlate with Hammett numbers. Symmetric dinitrile $\Delta\mu$ values decrease monotonically with increasing conjugation of the connecting bridge, likely due to improved mechanical coupling and, to a lesser extent, an increased population of inversion symmetric conformations.

Introduction

When a weak electric field is applied across a molecule, both the molecular vibrational energy levels and the transition dipoles between the levels change slightly. These changes affect the infrared absorption spectrum of the sample, and are collectively called the vibrational Stark effect (VSE). VSE measurements yield information on the physics of molecular vibrations, including the anharmonicity of vibrational modes and the perturbation of chemical bonds by electric fields. VSE measurements can also be used to calibrate the sensitivity of a vibrational frequency to an electric field, called the Stark tuning rate, which allows an environmentally-induced vibrational band shift to be used as a local probe of electrostatic fields^{1,2}. Different vibrational modes have widely varying Stark effects, so a VSE spectrum can sometimes be used to identify a weak vibrational band of interest from a collection of overlapping background bands².

Strong local electric fields are common in condensed matter systems, including the solvation of charged or dipolar molecules³, in ionic matrices, at electrode surfaces, across

biological membranes⁴, and in the organized environment of proteins^{5,6}. These local fields can be very large, typically lying in the range of 0.1 to 10 MV/cm. Vibrational band shifts have been studied in all these systems⁷⁻¹⁰, but in most cases neither the electric field nor the Stark tuning rate of the probe molecule was known accurately. We have recently shown that vibrational Stark spectroscopy is a useful tool for understanding vibrational band shifts for diatomic ligands bound to the heme iron in myoglobin^{1,2}, and it will likely prove useful for understanding and predicting band shifts in a wide variety of other systems.

In this work, a sensitive VSE measurement method has been developed and used to analyze the nitrile stretch mode of several aliphatic and aromatic nitriles and dinitriles. The nitrile stretch mode was chosen for analysis for several reasons. The frequency is high enough to be well separated from most other vibrational modes and is in a spectral region that is relatively easy to detect, as it occurs near the maximum detectivity of an indium antimonide infrared detector. Bands are generally intense and narrow, properties which contribute to a strong Stark effect. A normal mode analysis of organic nitriles shows that the nitrile stretch mode is highly localized to the nitrile bond¹¹, which both simplifies the interpretation of VSE data and implies that changes to the rest of the molecule minimally perturb the nitrile normal mode. Finally, there is a great deal of experimental and theoretical literature on nitrile vibrations¹²⁻¹⁸.

Theory

A vibrational Stark analysis requires an absorption spectrum of the sample, $A(\theta)$, and a Stark difference spectrum, $\Delta A_{\theta}(\theta)$, which is the absorption in the presence of an external field, minus the absorption without the field. θ is the angle between the light polarization vector and the direction of the applied electric field, and can be varied from 90 to about 45 degrees using the experimental setup described in the next section. As explained below, the absorption and Stark difference spectra are fit to give the Stark fit coefficients, from which are computed the Stark parameters of the sample.

Stark parameters: For an absorption band, Stark effects are expected to yield a slight peak shift for each molecule in a sample, without any lineshape change, where the shift is given by:

$$\Delta\nu = \frac{1}{hc} \left[\mu \cdot \mathbf{F} + \frac{1}{2} \mathbf{F} \cdot \alpha \cdot \mathbf{F} + \dots \right] \quad (1)$$

μ and α are called the difference dipole moment and the difference polarizability, respectively, referring to the differences between ground and excited states. These names are not entirely accurate for VSE due to field-induced bond changes^{19,20}, but the variables are still useful and completely general. An electric field also affects the transition dipole moment, $\mathbf{M}(\mathbf{F})$, through the transition polarizability, \mathbf{A} , and the transition hyperpolarizability, \mathbf{B} :

$$\mathbf{M}(\mathbf{F}) = \mathbf{M} + \mathbf{A} \cdot \mathbf{F} + \mathbf{F} \cdot \mathbf{B} \cdot \mathbf{F} + \dots \quad (2)$$

Again, the variables \mathbf{A} and \mathbf{B} are useful and completely general but the names are not entirely accurate for VSE.

As the samples studied in this work were immobilized in a frozen glass, the molecular orientations were isotropic (however, field-induced poling can be a problem and is discussed below). Integrating the peak shifts and extinction coefficient changes given in equations 1 and 2 over all orientations, combined with the sample absorption spectrum, yields a computed Stark spectrum. This computed spectrum is a function of the two field-independent parameters, ν and \mathbf{M} , and the four Stark parameters, $\Delta\nu$, $\Delta\mu$, \mathbf{A} , and \mathbf{B} .

Stark fit coefficients: With the assumption that the absorption lineshape is unaffected by the applied electric field, the computed Stark spectrum can be expressed as a sum of derivatives of the absorption spectrum²¹:

$$\Delta A_{\nu}(\nu) = |\mathbf{F}|^2 A_{\nu} A(\nu) + \frac{|\mathbf{F}|^2}{15hc} B_{\nu} \frac{\partial}{\partial \nu} \frac{A(\nu)}{\nu} + \frac{|\mathbf{F}|^2}{30h^2c^2} C_{\nu} \frac{\partial^2}{\partial \nu^2} \frac{A(\nu)}{\nu} \quad (3)$$

\mathbf{F} is the electric field vector and A_{ν} , B_{ν} , and C_{ν} , which are called the zeroth, first, and second derivative coefficients, respectively, are found from equations 1 and 2 to be:

$$A_{\nu} = \frac{1}{3|\mathbf{M}|^2} \text{Tr}(\mathbf{A}^T \mathbf{A}) + \frac{3\cos^2 \theta}{10|\mathbf{M}|^2} \left[\text{Tr}(\mathbf{A})^2 + \text{Tr}(\mathbf{A}^2) \right] \frac{2}{3} \text{Tr}(\mathbf{A}^T \mathbf{A}) + \frac{2}{3|\mathbf{M}|^2} M_i B_{ij} + \frac{3\cos^2 \theta}{5|\mathbf{M}|^2} M_i B_{jj} + M_i B_{jji} \frac{2}{3} M_i B_{ij} \quad (4)$$

$$B_{\square} = \frac{10}{|\mathbf{M}|^2} \mathbf{M} \cdot \mathbf{A} \cdot \square\square + \frac{3(3\cos^2\square\square - 1)}{|\mathbf{M}|^2} \square\square \cdot \mathbf{A} \cdot \mathbf{M} + \mathbf{M} \cdot \square\square \text{Tr} \mathbf{A} \square - \frac{2}{3} \mathbf{M} \cdot \mathbf{A} \cdot \square\square \square + \frac{5}{2} \text{Tr} \square\square + \frac{(3\cos^2\square\square - 1)}{|\mathbf{M}|^2} \square\square \frac{3}{2} \mathbf{M} \cdot \square\square \cdot \mathbf{M} \square - \frac{1}{2} |\mathbf{M}|^2 \text{Tr} \square\square \square \quad (5)$$

$$C_{\square} = 5|\square\square|^2 + (3\cos^2\square\square - 1)(3\cos^2\square\square - 1)|\square\square|^2 \quad (6)$$

In equation 6, \square is the angle between \mathbf{M} and $\Delta\square$. There are also third derivative and higher order terms in the derivative expansion, but they are all proportional to the fourth or higher powers of the field, a dependence which is expected to be small and which was not seen in the data. The derivative coefficients were found from experimental data by fitting a measured Stark difference spectrum with a sum of the frequency-weighted derivatives of the absorption spectrum using software developed for this purpose. The fitting procedure is described below.

As the \square dependence in equations 4-6 is restricted to a linear dependence on the term $3\cos^2\square - 1$, only 6 scalar values of the 48 tensor components of the Stark parameters can be determined from Stark measurements, so a number of assumptions are needed to simplify the problem. The transition hyperpolarizability, \mathbf{B} , is assumed to be negligible, as it has been calculated to contribute at most a few percent to A_{\square}^{22} . It is simplest to consider the molecular z axis to be parallel to the transition dipole moment, \mathbf{M} . In this system, $\Delta\square_{zz}$ and A_{zz} are parallel polarizability components, represented by $\Delta\square_{\parallel}$ and A_{\parallel} , and $\Delta\square_{xx}$, $\Delta\square_{yy}$, A_{xx} , and A_{yy} are the perpendicular polarizability components, represented by $\Delta\square_{\perp}$ and A_{\perp} , using the assumption that the perpendicular components are equal on the two axes. The off-diagonal components, representing polarizabilities perpendicular to an applied field, are assumed to be zero. With these simplifications, the six remaining Stark parameters are $|\Delta\square|$, $\Delta\square_{\parallel}$, A_{\parallel} , \square , $\Delta\square_{\perp}$, and A_{\perp} , which can be found from experimental data.

Further assumptions about the Stark parameters were made for data where the \square dependence of the Stark difference spectrum was not measured. Here, we assumed that a nitrile or dinitrile vibration behaves as a one-dimensional oscillator, with the result that \square , $\Delta\square_{\perp}$, and A_{\perp} are assumed to be zero. This assumption of a one-dimensional system is expected to be exact for nitriles with rotational symmetry about the nitrile bond (ignoring

solvent induced asymmetry); it was also shown to be reasonably good for samples where the \square dependence was measured, described below.

Due to the bulk dielectric properties of the solvent as well as specific field-induced perturbations of the solvent near the sample molecules, the electric field at a sample molecule is greater than the average electric field (total applied voltage divided by electrode spacing). This difference is corrected for using the local field correction factor f :

$$\mathbf{F}_{\text{local}} = f\mathbf{F}_{\text{external}} \quad (7)$$

Using the Lorentz spherical cavity model, which treats the solvent as a bulk dielectric, f is estimated to be between 1.1 and 1.3^{23,24}. While the exact value of f is not known, it is expected to be the same constant for all experiments in this work, as well as for a great deal of other work, allowing for meaningful comparisons. Due to the lack of knowledge of this factor, results are reported in terms of factors of f .

Experimental

Samples were solutions of small aliphatic and aromatic nitriles in the glass forming solvent, 2-methyl-tetrahydrofuran (2-MeTHF). VSE sample concentrations ranged from 0.1 M to 0.3 M, and no concentration dependent effects were observed throughout this range in either absorption or Stark spectra, indicating that nitrile dimers^{17,25} did not form to a measurable extent. In support of this assertion, room temperature absorption spectra of several nitriles (butyronitrile, hydrogen cyanide, valeronitrile, and deuterated acetonitrile) also showed no concentration dependent effects for concentrations between 2.4 mM and 0.8 M. Low temperature extinction coefficients were measured by taking spectra of known sample concentrations with a setup that was essentially identical to that used for Stark measurements. The 2-MeTHF solvent and all nitriles, except for hydrogen cyanide, were purchased from Aldrich and used without further purification.

For the synthesis of hydrogen cyanide (HCN), equal volumes of 2-MeTHF and a 0.3 M aqueous solution of potassium cyanide were put in a vial, forming an organic layer on top and an aqueous layer below. Dropwise addition of concentrated HCl formed the HCN product, which then partitioned into the organic layer from which the sample was

taken for spectroscopy. The partition coefficient (organic/aqueous) is 4.62, determined from room temperature infrared peak intensities of HCN in the aqueous layer in contact with various volumes of 2-MeTHF. Using the partition coefficient, a known concentration of HCN was dissolved in the organic layer and used to determine the low temperature extinction coefficient of HCN. It was shown with control experiments that no CN^- partitioned into the organic layer and that all measurable aqueous CN^- reacts to form HCN in the presence of excess acid. The room temperature peaks analyzed here are aqueous CN^- at 2080 cm^{-1} , aqueous HCN at 2093 cm^{-1} , and organic HCN at 2079 cm^{-1} .

The infrared spectroscopy cell consisted of a pair of 13 mm diameter by 1 mm thick calcium fluoride windows (Janos Technology Inc.) with approximately 50 \AA of nickel vacuum deposited on the surfaces facing the sample, a design which is similar to that used in previous work²⁶ and in electronic Stark spectroscopy²⁴. A layer of nickel typically transmits 50 to 80 percent of the incident light throughout the infrared region, with the rest being either absorbed or reflected, and has a resistance of about $5\text{ k}\Omega$ across the face of a window. The nickel electrodes were connected to a high voltage DC power supply (Trek Instruments Inc.), whose output voltage was synchronized to the FTIR scan timing, described below, with a home-built control unit. The windows were separated from each other by a pair of 28 micron thick Teflon spacers and held in place with a metal clamp. Using interference contour fringes, the windows were made parallel to within 1° to ensure both a uniform pathlength and a uniform field in the cell. The window separation in the center of the cell was measured to ± 0.05 micron after each experiment by taking a visible spectrum of the empty cell and measuring the frequency spacing between spectral interference fringes. After a sample was loaded into the cell by capillary uptake, the cell was immersed in liquid nitrogen to rapidly freeze the sample. This formed a transparent glass without observable cracks, crystallization, or other defects. The nitrogen immersion dewar is a custom built optical cryostat with several novel features designed to avoid bubbles and schlieren effects²⁷.

Spectroscopy was carried out on a Bruker IFS 66V/S FTIR, with a nitrogen cooled indium antimonide detector and 0.5 cm^{-1} resolution. We first tried to record Stark spectra with the FTIR in step-scan mode and by locking into the second harmonic of a 200 hertz

electric field applied to the sample. This is similar to the standard method for visible and near infrared Stark spectroscopy²⁴ and to a method used for measuring vibrational circular dichroism²⁸. However, the noise reduction achieved by lock-in detection was more than offset by the noise gained by using step-scan mode, rather than rapid scan mode. Step-scan methods proved to be difficult with a conventional lock-in amplifier because a large detector signal change is observed with each mirror step, requiring a delay of many lock-in time constants before a VSE signal can be observed. Better signal-to-noise was obtained using a DC electric field with the FTIR in its normal rapid scan mode. Using this method, a single scan is made with the field turned off, then a scan with 2.5 kilovolts applied to the sample in one direction (about 0.9 MV/cm electric field), then another field-off scan, and finally a 2.5 kilovolt potential in the opposite direction. The alternating field pattern was chosen to avoid charge build-up in the sample and to prevent sample poling; both field-on scans gave identical spectra and were averaged together. This cycle was repeated at least 512 times, which is about one hour, for each VSE measurement. At the end, the field-on and field-off interferograms were averaged and Fourier transformed, and the ratio of their spectra were used to compute a Stark spectrum. The field-off spectrum, relative to background scans with the sample out of the light path, provided a good quality absorption spectrum with exactly the same conditions as the Stark spectrum. Sample molecules did not change their orientations due to the applied electric field in the immersion cryostat to any measurable extent, as determined by (i) the lack of differences between positive and negative field results after a constant positive field had been applied for 15 minutes or longer and (ii) the lack of a difference between sequential field-on scans, using a different electric field pattern²⁹.

Data Analysis

Fits were made with at least one of three methods³⁰. Where the absorption spectrum did not have overlapping bands, the numerical derivatives of the measured spectrum were used so as to eliminate errors due to naturally asymmetric lineshapes or any consistent measurement error, such as light scatter or imperfect phase correction of the interferogram. This fitting method introduces the least error but cannot be used to determine separate Stark parameters for overlapping bands. For samples with

overlapping bands, and for several samples with single bands in order to check the reliability of the method, the absorption spectra were fit first with a modified Voigt lineshape for each peak. The form of a modified Voigt lineshape is a linear combination of a Lorentzian and a Gaussian, where the Lorentzian and Gaussian are constrained to have the same peak positions and the same full width at half maximum³¹. It is similar to a Voigt lineshape, a convolution of a Lorentzian and Gaussian, and has similar versatility, but is easier to fit. The analytical derivatives of the separate peaks were then used to fit the VSE data, providing separate fit coefficients for each peak. The third method used a simultaneous fit of the absorption spectrum with modified Voigt lineshapes, and of the Stark spectrum with the analytical derivatives of these lineshapes. The sum of the two fit errors was minimized, where each least-squares error was weighted by the respective spectral rms noise, in order to use all the spectral information available for all the fit parameters. The simultaneous fitting method is less sensitive to noise in the absorption spectrum, but more sensitive to noise in the Stark data. Where multiple methods were carried out on the same set of spectra, fit results were identical, within statistical error.

In order to achieve a good signal-to-noise ratio, most data were obtained only at the experimental angle \square at 90° , where unpolarized light can be used and less of the infrared beam is clipped by the sides of the sample. These data were interpreted using the one-dimensional assumptions, described above. Fits to these data constrained A_{\square} values to be non-negative, as negative values are impossible for a one-dimensional system and for a negligible transition hyperpolarizability (Eqn. 4). This constraint was only required for data sets of poorer quality, suggesting that those A_{\square} values were dominated by noise and that the constraint is appropriate.

In some situations, strong interference fringes were observed at normal incidence due to reflections between the sample cell windows, which were reduced by rotating the sample away from the normal by up to 20° (still using unpolarized light). Those data were corrected to normal incidence, again with the assumption that the nitrile Stark effects follow one-dimensional behavior. The corrections are relatively insensitive to the assumptions for small rotation angles and unpolarized light³².

Despite rotating some samples, Stark spectra often showed a non-zero baseline, typically dominated by spectral interference fringes (about 130 cm^{-1} period), but also sometimes including a uniform offset or slope. To prevent the baseline from affecting the fit parameters, the baseline was fit along with the rest of the Stark spectrum using a linear combination of a sine wave and a straight line. This did not interfere with the rest of the fitting because the baseline extended well beyond the nitrile peaks, and the functions used to fit the baseline yield very different curve shapes than the functions used to fit the VSE peaks.

The magnitudes of the transition dipoles were found from the area of the absorption band, expressed as an extinction coefficient spectrum, $\epsilon(\omega)$ ³³:

$$|\mathbf{M}| = \sqrt{\frac{3\epsilon_0 h c \ln 10}{2\pi^2 N_A \pi} \int \epsilon(\omega) d\omega} \quad (8)$$

Several samples showed multiple peaks in the nitrile stretch region. For situations where the peaks were determined to be due to multiple vibrational modes, the peak areas were integrated separately using fits with modified Voigt lineshapes, as described above (these samples include all dinitriles with multiple peaks and deuterated acetonitrile). On the other hand, where multiple peaks were determined to be due to either heterogeneity or Fermi resonance, the integration was carried out over all the peaks in order to include the entire sample and all the intensity of the nitrile mode (these samples include HCN and other aliphatic nitriles). Implicit in the latter method is the assumption that the transition dipole moments have the same magnitude for all nitrile stretches in a heterogeneous sample.

Results

Aliphatic nitriles: The nitrile stretch mode of acetonitrile gives rise to an intense and narrow absorption band (Fig. 1A). There is also a weak Fermi resonant peak 50 cm^{-1} to higher energy (not shown; peak maximum is at 2297.1 cm^{-1} , intensity is $6.90\text{ M}^{-1}\text{cm}^{-1}$, FWHM is 20.1 cm^{-1}), which arises from resonance between the nitrile mode and the combination mode of the C-C stretch and the symmetric C-H bend⁸. To facilitate comparison with other spectra, Figure 1 and similar figures in this paper are scaled to




Figure 1. Spectra of acetonitrile. These and subsequent spectra are of samples dissolved in 2-MeTHF, at 74 K, and at normal incidence to the infrared beam. (A) Extinction coefficient spectrum of the nitrile stretch mode; the line is not a fit, but is simply shown to guide the eye. (B) Dots represent the Stark data, the solid line is the best fit to the Stark spectrum, using numerical derivatives of the data in panel A, and the dashed line is the Stark baseline. In this and subsequent figures, Stark spectra are scaled for a 1 MV/cm applied electric field. (C) The solid line is the fit from panel B, with the baseline subtracted, and the dashed lines are the zeroth, first, and second derivative contributions.

show spectral extinction coefficients and to show Stark spectra for an applied field of 1 MV/cm (Eqn. 3). The unscaled acetonitrile data, which are fairly typical, had a maximum absorption of 0.21 O.D., a minimum Stark absorption, relative to the baseline, of -4.4×10^{-4} O.D. using a 0.88 MV/cm electric field, and Stark rms noise levels of 5.2×10^{-6} O.D.

Table 1. Vibrational Stark effect data for unconjugated aliphatic nitriles.

Compound	Absorption data				Stark data					
	$\bar{\nu}^a$	peak area	$ M $	q	$A_{\bar{\nu}}$ 10^{20}	$B_{\bar{\nu}}$ 10^{40}	$C_{\bar{\nu}}$ 10^{63}	$ \Delta\bar{\nu} ^b$	$\Delta\bar{\nu}_{ }^{b,c}$	$A_{ }^{b,c}$
	cm^{-1}	$\text{M}^{-1}\text{cm}^{-2}$	D	e	m^2/V^2	Jm^2/V^2	$\text{J}^2\text{m}^2/\text{V}^2$	D/f	$\text{\AA}^3/f^2$	$\text{\AA}^3/f$
HCN L^d	2054.9	3989	0.134	0.79	15.8	2.23	28.89	0.0588	-4.93	3.55
HCN M^d	2064.7	3989	0.134	0.79	0	-2.92	20.12	0.0491	-1.76	0
HCN H^d	2072.4	3989	0.134	0.79	0	0.43	14.89	0.0422	0.26	0
acetonitrile ^e	2247.5	830	0.058	0.36	1.8	0.22	5.56	0.0258	-0.81	0.53
propionitrile	2243.5	778	0.056	0.35	1.9	0.52	6.42	0.0277	-0.71	0.52
butyronitrile ^e	2247.0	813	0.058	0.35	1.9	0.92	6.69	0.0283	-0.48	0.53
valeronitrile ^e	2239.2	1024	0.065	0.40	0.5	0.87	5.95	0.0267	0.14	0.23
hexanenitrile ^e	2243.2	793	0.057	0.35	0.5	-0.03	6.55	0.0280	-0.52	0.25
acetonitrile-d3	2255.5	974	0.063	0.37	2.1	-0.17	6.30	0.0275	-1.14	1.09
acetonitrile-d3 (C-D) ^f	2234.5	904	0.061	0.39	7.2	1.77	14.16	0.0412	-1.89	0.60

a) Peak maximum. b) Assumes one-dimensional behavior, *i.e.* $\bar{\nu}=0^\circ$, and off-diagonal and perpendicular components of $\Delta\bar{\nu}$ and \mathbf{A} are 0. c) \mathbf{B} is assumed to be 0. d) Peak areas for HCN low (L), medium (M), and high (H) energy nitrile absorptions are combined. e) Fermi resonant peaks are observed, and their areas are combined with the nitrile peak area. f) C-D stretch mode of deuterated acetonitrile.

As seen in Figure 1C, the VSE spectrum is dominated by the second derivative of the absorption spectrum, but also shows significant zeroth and first derivative components (Table 1), all of which are highly reproducible. The $\bar{\nu}$ angle dependence of the Stark spectrum was studied as well and is discussed below; however, it is ignored here and in Table 1, using the one-dimensional assumptions, to allow for a meaningful comparison with other nitriles where the $\bar{\nu}$ dependence was not measured. From the second and zeroth derivative contributions, respectively, $|\Delta\bar{\nu}|$ is 0.0258 ± 0.0004 D/f and

A_{\parallel} is $0.53 \pm 0.03 \text{ \AA}^3/f$, which imply a 0.43 cm^{-1} shift and a 6.1% increase of band intensity for molecules aligned parallel to a 1 MV/cm electric field (errors represent 1 standard deviation for 5 data sets). As A_{\parallel} is more than sufficient to account for the entire first derivative contribution, $\Delta\epsilon_{\parallel}$ is found to be slightly negative, with a value of $-0.81 \pm 0.07 \text{ \AA}^3/f^2$, which implies a band shift of 0.02 cm^{-1} to higher energy for molecules aligned parallel to a 1 MV/cm field. Due to the large width and low intensity of the resonant band, as well as its overlap with infrared absorption from liquid nitrogen in the immersion cryostat^{27,34}, its Stark effect could not be measured reliably. However, its $|\Delta\epsilon_{\parallel}|$ value appears to be between 0.01 and 0.05 D/f, putting it in the same range as the nitrile band Stark effect.

Deuterating acetonitrile lowers the C-H frequencies, which moves the Fermi resonant band well below the nitrile mode and moves the C-H stretch frequency to just below the nitrile mode (Fig. 2A,B). The two peaks were determined to have different Stark characteristics by the fact that derivatives of the entire absorption spectrum cannot be satisfactorily fit to the Stark spectrum. This required a separation of the peaks to allow independent fitting. However, the absorption spectrum also cannot be fit with two modified Voigt lineshapes due to the relatively large absorption between the peaks. This extra absorption indicates the presence of non-linear coupling between the C-N and C-D stretching modes³⁵. A theory has not been developed yet for Stark effects of resonant modes so an interpolation method was used to separate the peaks. This approach yielded a good fit to the data despite its lack of a physical basis. Two modified Voigt lineshapes, called Fit_L and Fit_H , were fit to the low and high energy peaks of the absorption spectrum, respectively, while ignoring the fit error of the region between the peaks (2237 to 2250 cm^{-1}). The ratios of the fits were used to separate the experimental spectrum, A , yielding separate spectra for the two peaks, A_L and A_H (these are all functions of ϵ):

$$A_L = \frac{A \cdot Fit_L}{Fit_L + Fit_H} \quad A_H = \frac{A \cdot Fit_H}{Fit_L + Fit_H} \quad (9)$$

As $A_L + A_H = A$, all the sample absorption is included without requiring the use of additional fitted peaks; this interpolation method also preserves the natural peak asymmetry. Derivatives of A_L and A_H were then fit to the Stark data (Fig. 2B). The good

quality of the fit away from the region between the bands suggests that the fit parameters are likely to be representative of the isolated normal modes, and are thus comparable to parameters for other nitriles. Not surprisingly, the nitrile VSE parameters for deuterated acetonitrile (Table 1) are similar to those for protonated acetonitrile. For the C-D stretch mode, $|\Delta\mu|$ is $0.0412 D/f$, which is about 50% larger than the nitrile difference dipole.




Figure 2. Spectra of CD_3CN , butyronitrile, and HCN . (A) Dots show the extinction coefficient spectrum for CD_3CN , showing a C-D stretch mode at 2235 cm^{-1} and the C-N stretch mode at 2256 cm^{-1} . The solid line is a sum of two modified Voigt lineshapes, demonstrating the resonant absorption between the bands; these lineshapes were used to separate the measured absorbance into two bands (see text). (B) Dots represent the CD_3CN Stark spectrum, the solid line shows the best fit to the Stark data, using different parameters for the two bands, and the dashed line shows the Stark baseline. (C,D) Extinction coefficient and Stark spectra for butyronitrile, where the two peaks are interpreted as being due to Fermi resonance. The line in panel C is shown to guide the eye, whereas the line in panel D is a fit to the Stark spectrum using numerical derivatives of the absorption spectrum, fit with a single set of coefficients. (E,F) Extinction coefficient and Stark spectra for HCN , where the three peaks are interpreted to result from heterogeneous hydrogen bonding to the solvent. Solid lines show fits that were made simultaneously to the two spectra, using modified Voigts for the extinction coefficient spectrum and their analytical derivatives for the Stark spectrum. Dashed lines in panel E show the three modified Voigt lineshapes used to fit the peak, and the dashed line in panel F is the Stark baseline.

A homologous series of aliphatic nitriles was studied, from hydrogen cyanide to hexanenitrile. With the exception of HCN (discussed below), the transition dipoles and difference dipoles are nearly identical across the series (Table 1). Other Stark parameters vary some, but with no clear trends and with less magnitude than the variation between aliphatic and other nitriles. The molecular polarizability changes by about a factor of 3 from acetonitrile to hexanenitrile³⁶, suggesting that the molecular polarizability has a minimal influence on vibrational Stark effects for nitriles.

Lengthening the hydrocarbon chain to two units in propionitrile results in an absorption spectrum with a single peak and no discernable Fermi resonance. However, on adding one, two, or three more CH₂ units to the hydrocarbon chain, giving butyronitrile, valeronitrile and hexanenitrile, respectively, the absorption spectra show a pair of peaks (Figure 2C shows butyronitrile). In all cases, the peaks fit well with a pair of modified Voigt lineshapes, without any extra absorption between the peaks. Using these fits, the separations between the peaks are 9.7, 13.0, and 13.0 cm⁻¹ for the respective nitriles, while the ratios of the areas of the lower and higher frequency peaks are 0.22:1, 1.38:1, and 0.006:1 (all at 74K). Possible assignments for the multiple peaks include Fermi resonant interactions as seen for acetonitrile, heterogeneous sample conformations involving *anti* and *gauche* conformations, partial dimer formation in the sample, solvent heterogeneity, or a combination of these factors. These assignments don't affect the analysis of VSE data, although the interpretation varies considerably. From a variety of experiments, all the possible assignments involving simple heterogeneity can be eliminated: (i) spectra are independent of sample concentration over a wide range of concentrations at both room temperature and at 74K, indicating that dimer formation does not occur at the concentrations used; (ii) the relative areas of the peaks are different for different compounds, and none agree with the Boltzmann population calculations that the ratio of *gauche:anti* populations are expected to be about 0.07:1 in the frozen samples or 0.6:1 at room temperature³⁷; (iii) the peak separations are much larger than those between homologous nitriles or than those calculated with AM1 methods for *gauche* and *anti* conformations (<1 cm⁻¹), also arguing against assignment based on conformational heterogeneity; (iv) room temperature spectra show multiple peaks and the solvent phase transition is not apparent in temperature dependent spectra,

suggesting that solvation is not primarily responsible for the multiple peaks; and (v) room temperature spectra in toluene are very similar to those in 2-MeTHF, indicating again that specific solvation effects are not important. Thus, simple heterogeneity is unlikely, and the multiple peaks are almost certainly due, at least in part, to Fermi resonance. However, it remains surprising that Fermi resonance would yield significantly different spectra for each of the aliphatic nitriles studied. It is also surprising that valeronitrile shows a strong temperature dependence, whereas the other nitriles show a weak temperature dependence.

As the intensity of the multiple peaks for butyronitrile, valeronitrile, and hexanenitrile is expected to arise from the nitrile mode (regardless of assignment), the entire bandshapes were integrated to find the transition dipole moments. From VSE spectra (Fig. 1D shows butyronitrile), it was found that the Stark parameters of the pairs of peaks are identical within measurement error, so they are reported in Table 1 with one value for each nitrile. Assuming the Fermi resonance assignment is correct, it is found that the Stark parameters of these Fermi resonant bands are identical to those of the parent bands, suggesting that VSE parameters represent the Stark effects of the source of the absorption intensity, rather than those of the forbidden mode. In contrast, recent work on the VSE of nitric oxide bound to the heme iron in myoglobin showed a case where Fermi resonance affects the Stark effect of the parent band². In general, Stark effects of Fermi resonances are not well understood.

The simplest member of the aliphatic series, hydrogen cyanide, does not behave like the other aliphatic nitriles. Its absorption band is at lower energy, is broader, and is about 4 times more intense (Fig. 2E,F). The simplest fit that can be made to the absorption and Stark spectra requires three overlapping modified Voigt lineshapes, characterized by different Stark parameters. Due to the simplicity of HCN, there are no resonant modes or alternate conformations; consequently, the three peaks most likely represent heterogeneous hydrogen bonding from the sample to the 2-MeTHF solvent. This interaction, which cannot occur for the other samples, would be expected to significantly affect the nitrile bond. In agreement with this assignment, the spectrum of HCN exhibits a single broad peak when it's dissolved in 2-MeTHF at room temperature,

and a single sharp peak when in a toluene glass at low temperature. The transition dipoles were considered to be the same for the three heterogeneous peaks, despite the fact that this is likely to be a poor assumption, because they cannot be determined independently. It was found that both the transition dipole moment and the difference dipole moments of HCN are much larger than they are for the other aliphatic nitriles.

Aromatic nitriles: With a few exceptions, aromatic nitrile absorption spectra show a single strong band in the nitrile stretch region. The spectrum of 4-chloro-benzonitrile is shown in Figure 3 and its Stark parameters are listed in Table 2 along with those of other aromatic nitriles. A notable exception is the symmetric compound 1,4-dicyanobenzene, which is discussed below with the other symmetric dinitriles. Also, benzonitrile and 2-chloro-benzonitrile show weak absorptions at 16 and 35 cm^{-1} higher frequency, respectively, than the strong nitrile band. The Stark spectra of the small peaks fit well with the same derivative components as the major peaks, but were too small for a thorough analysis.

As with the aliphatic nitriles, fits to the aromatic Stark spectra are dominated by the second derivative contribution, and the source of the Stark effect is dominated by the difference dipole moment. However, aromatic nitrile VSE spectra exhibit slightly larger zeroth derivative contributions and much larger first derivative contributions, leading to significantly different lineshapes. Upon analysis, it is seen that the large first derivative contribution is due to larger difference dipoles and larger transition polarizabilities, rather than to a significant change in difference polarizabilities (Table 2), which are again small and slightly negative. This result contrasts with our earlier work on 4-methoxy-benzonitrile in which it was concluded that the zeroth derivative contribution was sufficiently small to be ignored, and that the entire first derivative contribution represented a positive difference polarizability²⁶. The change in interpretation is appropriate, because of the substantial improvement of data quality that has been achieved.

In contrast to the aliphatic nitrile data, aromatic nitrile VSE spectra show a relatively large variation in $|\Delta\mu|$ values with molecular structure. As seen in Figure 4, it




Figure 3. Spectra of 4-chloro-benzonitrile. (A) Extinction coefficient spectrum, where the line is shown to guide the eye. (B) Dots represent the Stark spectrum, the solid line is the best fit to the Stark spectrum using numerical derivatives from panel A, and the dashed line is the Stark baseline. (C) The solid line is the fit from panel B with the baseline subtracted, and the dashed lines are the zeroth, first, and second derivative contributions.

is found that both the transition dipole and difference dipole moments correlate well with Hammett parameters, which are unitless values that characterize the electron donating or electron withdrawing nature of substituents on an aromatic ring³⁸. As the transition

Table 2. Vibrational Stark effect data for aromatic nitriles.

Compound	Absorption data					Stark data				
	$\bar{\nu}^a$	peak area	$ M $	q	A_{\parallel}^b	B_{\parallel}^b	C_{\parallel}^b	$ \Delta\mu ^{b,c}$	$\Delta\mu_{\parallel}^{b,c}$	$A_{\parallel}^{b,c}$
	cm^{-1}	$\text{M}^{-1}\text{cm}^{-2}$	D	e	m^2/V^2	Jm^2/V^2	$\text{J}^2\text{m}^2/\text{V}^2$	D/f	$\text{\AA}^3/f^2$	$\text{\AA}^3/f$
benzonitrile	2227.8	2083	0.093	0.57	2.6	1.52	10.86	0.0361	-0.65	1.00
1,2-dicyanobenzene ^d	2235.3	819	0.058	0.36	1.4	2.01	5.80	0.0263	0.60	0.32
1,3-dicyanobenzene ^d	2236.3	868	0.060	0.37	5.5	2.23	9.08	0.0330	-0.71	0.92
1,4-dicyanobenzene (A) ^{d,e}	2231.7	862	0.060	0.37	2.1	2.76	3.20	0.0196	0.89	0.58
1,4-dicyanobenzene (S) ^{d,f}	2243.7	138	0.024	0.15	4.9	1.95	1.30	0.0125	0.43	0.35
2-Cl-benzonitrile	2232.1	1205	0.070	0.43	3.4	1.94	7.81	0.0306	-0.35	0.87
3-Cl-benzonitrile	2232.7	1260	0.072	0.44	2.9	2.04	8.45	0.0318	-0.23	0.82
4-Cl-benzonitrile	2230.6	1921	0.089	0.55	2.7	2.10	10.10	0.0348	-0.26	0.97
4-methoxybenzonitrile	2223.5	3724	0.124	0.76	1.8	1.33	20.60	0.0497	-0.02	0.49

a,b, and c) See the corresponding notes in Table 1. d) peak area and transition dipole are divided by two, to give results on a per nitrile basis. e) Antisymmetric (A) nitrile mode. f) Symmetric (S) nitrile mode.

dipole of a vibrational excitation is directly related to the partial charges on the excited group, this correlation with Hammett parameters is reasonable. The observation that the difference dipoles also correlate with Hammett parameters suggests that $|\Delta\mu|$ is determined by the same partial charges, a result which is quantified below and is discussed in a separate publication²⁰.

Dinitriles: Centrosymmetric compounds are particularly interesting for Stark analysis because all parameters that are linear with the electric field should vanish by symmetry, including $\Delta\mu$ and \mathbf{A} (eqns. 1 and 2). Also, an inversion symmetric compound has both symmetric and antisymmetric normal modes, where the former are expected to have no absorption intensity and the latter are expected to have twice the intensity observed for comparable non-symmetric compounds. The symmetric modes would be expected to gain intensity in an electric field due to the symmetry breaking of the field and the effect of the transition hyperpolarizability. Upon a weakening of the coupling between the two nitrile groups of a symmetric dinitrile relative to coupling to other internal degrees of freedom and to the solvent, the nitrile vibrations become more

independent. Thus, absorption and Stark spectra are expected to approach those of non-symmetric molecules.




Figure 4. Correlation of the transition dipole and difference dipole moments for the aromatic mononitriles and non-symmetric aromatic dinitriles with Hammett numbers. 4-OCH₃ is 4-methoxybenzonitrile; C₆H₅CN is benzonitrile; 2-Cl, 3-Cl, and 4-Cl are 2-, 3-, and 4-chlorobenzonitrile; and 1,2-CN and 1,3-CN are 1,2- and 1,3-dicyanobenzene.

As expected, most dinitriles show two absorption peaks in the nitrile stretch region. With the exception of *cis,cis*-mucononitrile (1,4-dicyano-*cis,cis*-1,3-butadiene), one peak is always about an order of magnitude stronger than the other, giving the antisymmetric and symmetric mode assignments, from the discussion above. 1,4-dicyanobenzene (not shown) exhibits strong resonance effects between the two nitrile modes, leading to an absorption spectrum analogous to that of deuterated acetonitrile (Fig. 2A,B), and requiring a similar analysis method (Eqn. 9).

Table 3. Vibrational Stark effect data for aliphatic dinitriles.

Compound	Absorption data				Stark data					
	$\bar{\nu}^a$	peak area ^d	$ M ^d$	q	$A_{\bar{\nu}}^e$ 10^{20}	$B_{\bar{\nu}}^e$ 10^{40}	$C_{\bar{\nu}}^e$ 10^{63}	$ \Delta\bar{\nu} ^b$	$\Delta\bar{\nu}_{ }^{b,c}$	$A_{ }^{b,c}$
	cm^{-1}	$\text{M}^{-1}\text{cm}^{-2}$	D	e	m^2/V^2	Jm^2/V^2	$\text{J}^2\text{m}^2/\text{V}^2$	D/f	$\text{\AA}^3/f^2$	$\text{\AA}^3/f$
succinonitrile	2249.9	374	0.039	0.24	4.7	0.04	6.42	0.0277	-1.58	0.57
adiponitrile (A) ^e	2244.4	697	0.053	0.33	0	0.70	5.75	0.0262	0.42	0
adiponitrile (S) ^f	2261.4	68	0.017	0.10	0	4.74	6.28	0.0274	2.84	0
fumaronitrile (S) ^f	2225.0	23	0.010	0.06	192.0	4.87	4.71	0.0237	-5.91	0.91
fumaronitrile (A) ^e	2236.5	268	0.033	0.20	0	0.32	0.71	0.0092	0.19	0
1,4-dicyano-2-butene	2251.3	515	0.046	0.28	0.2	1.24	2.84	0.0183	0.62	0.09
<i>cis,cis</i> -mucononitrile (L) ^g	2214.7	587	0.049	0.30	9.7	3.15	2.45	0.0171	0.46	1.03
<i>cis,cis</i> -mucononitrile (H) ^h	2223.1	548	0.048	0.29	0.9	-0.16	2.38	0.0168	-0.38	0.21

a,b, and c) See the corresponding notes in Table 1. d) Peak areas and transition dipoles are divided by two to give results on a per nitrile basis. e) Antisymmetric (A) nitrile mode. f) Symmetric (S) nitrile mode. g) Low (L) energy peak. h) High (H) energy peak.

The VSE predictions are borne out in the symmetric dinitrile data, presented in Table 3 and Figure 5, although other factors partially obscure this simple picture. In particular, the symmetric and anti-symmetric mode frequencies are close enough for dinitriles that there is often significant intensity sharing between the bands due to non-linear coupling³⁵. Also, it is calculated that approximately 7% of succinonitrile (dicyanoethane) and 12% of adiponitrile (1,4-dicyanobutane) molecules are expected to be in low symmetry conformations³⁷. Additionally, the dinitrile Stark data often exhibited a poor signal-to-noise ratio due to low intensity absorptions, small Stark effects, and low dinitrile solubilities in 2-MeTHF. Cyanogen³⁹ is likely to be among the most interesting compounds due to its high symmetry and the strong coupling between the nitriles, but was found to be insoluble and thus could not be analyzed.

While significant linear Stark effects were seen for all dinitriles studied, the difference dipole moments for both bands were invariably lower for dinitriles than for the comparable mononitriles. Furthermore, for the aliphatic dinitriles, the $|\Delta\bar{\nu}|$ values for both bands decreased with increased conjugation of the hydrocarbon bridge. This trend (Fig. 5) starts with succinonitrile and adiponitrile (Fig. 5A,B), which have $|\Delta\bar{\nu}|$ values and Stark spectra that are similar to those of aliphatic mononitriles. Increasing the

succinonitrile conjugation to give fumaronitrile (dicyanoethene) and the adiponitrile conjugation to give 1,4-dicyano-2-butene (Fig. 5C,D) and *cis,cis*-mucononitrile (Fig. 5E,F), yield significantly different Stark spectra. These spectra reveal progressively smaller second derivative components, leading to smaller difference dipoles. This behavior is expected from the discussion above because the mechanical coupling between the nitriles increases with increasing conjugation. The presence of low symmetry conformations may contribute to the observed trend, but this contribution is expected to be small based on the conformational calculations presented above.




Figure 5. Spectra of a homologous series of dinitriles. (A, B) Extinction coefficient and Stark spectra of adiponitrile (1,4-dicyanobutane); peaks at 2244 and 2261 cm^{-1} represent the antisymmetric and symmetric modes, respectively. The solid line in panel A is a fit of modified Voigts, which was used to separate the measured absorption into two bands. (C, D) Extinction coefficient and Stark spectra of 1,4-dicyano-2-butene. The VSE spectrum is fit with the numerical derivatives of the absorption. (E, F) Extinction coefficient and Stark spectra of *cis,cis*-mucononitrile (1,4-dicyano-*cis,cis*-1,3-butadiene). The VSE spectrum is fit with analytical derivatives of two modified Voigt lineshapes, fit to the absorption in panel E.

Angle dependence: After a survey of the Stark effects of a diverse collection of nitriles, acetonitrile (Fig. 1) and 4-chloro-benzonitrile (Fig. 3) were chosen for more detailed analysis due to their single nitrile peaks, minimal resonant interactions, high extinction coefficients, and high molecular symmetry. In order to examine the one-




Figure 6. Polarization angle dependence of the fit coefficients for acetonitrile (circles) and 4-chlorobenzonitrile (crosses). Error bars represent one standard deviation. Dashed lines are drawn with a slope-to-intercept ratio of 0.4 as guides to show the expected behavior for fit coefficients of one dimensional systems (see text).

dimensional assumptions made above, the zeroth, first, and second derivative fit contributions were plotted as a function of the angle parameter, $1-3\cos^2\theta$ (Fig. 6). For a purely one dimensional system ($\theta=0^\circ$ and off-diagonal and perpendicular components of $\Delta\alpha$ and \mathbf{A} are zero), the theoretical slope to intercept ratio for all three derivative contributions is 0.4, using equations 4-6.

It was found that 4-chloro-benzonitrile exhibits nearly one dimensional behavior for all three derivative components, with $\theta=5^\circ$, $A_\perp/A_\parallel=0.11$ and $\Delta\alpha_\perp/\Delta\alpha_\parallel=-0.001$. Using the fits to the angle dependent data, $|\Delta\alpha|$ is $0.0348 \text{ D}/f$, A_\parallel is $1.09 \text{ \AA}^3/f$, and $\Delta\alpha_\parallel$ is $-0.20 \text{ \AA}^3/f^2$, which are in close agreement with the values calculated with the one-dimensional approximations (Table 2).

Acetonitrile only exhibits one dimensional behavior for the second derivative component, showing much steeper slopes for the other two components. Analysis shows that $\cos\theta=1.04 \pm 0.002$ (clearly leading to an impossible θ value), $A_\perp/A_\parallel=0.39$, $\Delta\alpha_\perp/\Delta\alpha_\parallel=0.043$, $|\Delta\alpha|=0.0263 \text{ D}/f$, $A_\parallel=0.79 \text{ \AA}^3/f$, and $\Delta\alpha_\parallel=-0.60 \text{ \AA}^3/f^2$. The latter three parameters, $|\Delta\alpha|$, A_\parallel , and $\Delta\alpha_\parallel$, are in moderate agreement with the values calculated with the one-dimensional assumptions (Table 1). From the polarizability ratios, A_\perp/A_\parallel and $\Delta\alpha_\perp/\Delta\alpha_\parallel$, the dominant cause of both the zeroth and first derivative slopes would be interpreted to be a significant transition polarizability perpendicular to the bond axis. However, due to the three-fold rotational symmetry of acetonitrile, a perpendicular component (or an off-diagonal component) of the transition polarizability cannot occur without significant symmetry breaking by asymmetric solvation. A more likely explanation is that the homogeneous band shape of the nitrile mode changes in an electric field, most likely due to the weak Fermi resonant band 50 cm^{-1} to higher energy. This possibility is explicitly ignored in the Stark analysis presented here, likely leading to a mis-interpretation of the A_θ , B_θ , and C_θ angle dependence. This explanation is supported by the observation that statistically significant curvature is seen in Figure 6B for acetonitrile and that $\cos\theta > 1$, neither of which can be explained by the theory.

Discussion

It is found that difference dipoles correlate well with transition dipoles for all the mononitriles (Fig. 7), where the slope of a linear fit is $0.33/f$ and the intercept is 0.007

D/f . Dinitrile $|\Delta\mu|$ values generally agree with the correlation as well, but with significantly more scatter. In agreement with the predictions above, symmetric dinitriles exhibit small difference dipoles for both symmetric and antisymmetric modes and small transition dipoles for symmetric modes. The decrease of transition dipoles for antisymmetric modes (computed on a per nitrile basis) is expected for electrostatic reasons since two electron withdrawing groups compete for a finite amount of polarizable charge, thus decreasing the amount of partial charge on each nitrogen.




Figure 7. Correlation of difference dipoles with transition dipoles for all samples. Solid circles represent aliphatic mononitriles, solid squares represent aromatic mononitriles, open circles represent aliphatic dinitriles, and open squares represent aromatic dinitriles. The solid line is the best fit to the aliphatic and aromatic mononitrile data, with a slope of $0.33/f$ and intercept of $0.007 D/f$.

The correlation for mononitriles can be interpreted with a very simple model, which is presented elsewhere in more detail^{19,20,22}. The transition dipole of an harmonic oscillator is given by

$$|M| = \frac{q}{2\nu} \sqrt{\frac{h}{2cm\nu}} \quad (10)$$

where q is the effective charge of the oscillator and m is the reduced mass. From AM1 calculations on the mononitriles studied here, reduced masses are found to be 6.55 ± 0.05 atomic mass units. The small range reflects the fact that nitrile vibrations are highly

localized to the nitrile group. As the vibrational frequency is similar for all the mononitriles as well, the variation of the transition dipoles reflects, for the most part, the variation of the effective charge.

A simple interpretation of the difference dipole is that it reflects a displacement of the effective charge by a distance Δx , where the displacement is due to the anharmonicity of the potential surface:

$$|\Delta\mu| = q\Delta x \quad (11)$$

From equations 10 and 11, the slope of the correlation shown in Figure 7 (with slight corrections to account for the different μ values) corresponds to a Δx value of $0.012 \text{ \AA}/f$. The physical interpretation is that the expectation value of the C-N bond length is $0.012 \text{ \AA}/f$ longer in the first vibrationally excited state than it is in the ground state. This displacement can be compared against that expected from published values of the anharmonicity. In a perturbed harmonic oscillator model⁴⁰, the potential is defined as:

$$U(x) = \frac{k}{2}x^2 + \frac{\mu_3}{6}x^3 \quad (12)$$

The cubic anharmonicity of the C-N bond, μ_3 , has been determined from HCN vapor phase spectroscopic data⁴¹ to be 126 aJ/\AA^3 , leading to a Δx value of 0.0078 \AA . Thus, about 70% of the difference dipole values may be explained as being due to nitrile bond anharmonicity (assuming an f value of about 1.1). The discrepancy between the Δx values may result from: *i*) factors other than anharmonicity that contribute to $|\Delta\mu|$ values, *ii*) differences between the nitrile bond of HCN and of normal modes of larger nitriles, and *iii*) solvation effects on bond anharmonicity.

In Table 4, hydrogen cyanide and acetonitrile VSE results are compared against theoretical calculations of the molecules in the absence of solvent²². It is seen that all the experimental Stark parameters are several times greater than the calculated values, and are often larger by more than an order of magnitude. However, consideration of the Stark parameters in relation to the transition dipole values results in significantly better agreement. In particular, calculations for the acetonitrile transition polarizability and the difference dipole moment are within about 25% of the measured values. The large

discrepancies seen for hydrogen cyanide are not surprising due to the strong solvent effects that were observed.

Table 4. Calculated and observed vibrational Stark effects of HCN and CH₃CN.

property	unit	HCN		CH ₃ CN	
		obs ^a	calc ^b	obs ^c	calc ^b
M	D	0.134	0.0055	0.0582	0.0172
$\Delta\alpha$	D	0.05/ <i>f</i>	0.0010	0.0263/ <i>f</i>	0.0066
$\Delta\alpha_{ }$	Å ³	-2/ <i>f</i> ²	0.0402	-0.60/ <i>f</i> ²	0.0425
A	Å ³	1.2/ <i>f</i>	0.157	0.79/ <i>f</i>	0.171
$\Delta\alpha$ / M	unitless	0.37/ <i>f</i>	0.182	0.452/ <i>f</i>	0.384
$\Delta\alpha_{ }$ / M	(MV/cm) ⁻¹	-0.05/ <i>f</i> ²	0.0244	-0.034/ <i>f</i> ²	0.0082
A / M	(MV/cm) ⁻¹	0.03/ <i>f</i>	0.0952	0.045/ <i>f</i>	0.033

a) Unweighted averages of the values for the three sub-bands; assumes one dimensional behavior and that **B** is 0. b) Highest level of calculation from Reimers and Hush²². c) From data where α angle dependence was measured; values do not assume one dimensional behavior, but do assume **B** is 0.

Conclusions

A method for measuring vibrational Stark spectra has been developed and applied to the infrared absorption of the nitrile stretch mode of several aliphatic and aromatic nitriles. Most compounds were not analyzed with enough repeated experiments for detailed error estimates. Nevertheless, the precision of the method can be reasonably estimated by averaging the standard deviations for all repeated experiments. Difference dipoles, which invariably dominated the Stark effects, were reproducible to within ± 0.001 D/*f*, giving a Stark tuning precision of $\pm 0.02/f$ cm⁻¹/(MV/cm). Difference polarizabilities and transition polarizabilities led to smaller absorption changes, and could be measured reliably with precisions of 0.3 Å³/*f*² and 0.1 Å³/*f*, respectively.

The VSE data show that | $\Delta\alpha$ | values are typically smallest for symmetric dinitriles, larger for aliphatic nitriles and largest for aromatic species. The values of | $\Delta\alpha$ | are virtually all within the range of 0.01 D/*f* to 0.04 D/*f*, which is several orders of magnitude smaller than typical values for electronic Stark effects²⁴ and a factor of 2 to 8

smaller than those seen for the VSE of carbon monoxide¹ or nitric oxide² bound to the heme iron in myoglobin and heme model compounds. Where comparable experimental data exist, the $|\Delta\mu|$ values are in good agreement^{9,26}. Compared to theoretical results, they are several times larger than those predicted from *ab initio* calculations for isolated molecules²², but agreement improves significantly when Stark parameters are considered relative to transition dipoles.

While the nitrile stretch mode is typically well separated from other molecular vibrations, only about half of the samples studied exhibited well defined single bands. The other samples had multiple bands arising from Fermi resonant interactions, varying degrees of hydrogen bonding, and perhaps multiple sample conformations. Most dinitriles exhibited multiple bands as well, reflecting the symmetric and antisymmetric stretch modes. It was found that Stark parameters were generally similar for bands that arose from Fermi resonance, whereas they differed between bands arising from different modes. While the dinitrile data were not as simple as their symmetry might suggest, both Stark and absorption results agreed with expected trends, showing small difference dipoles and greater symmetry effects upon improved mechanical coupling.

For the aromatic nitriles studied, Hammett numbers correlate with both transition dipole moments and $|\Delta\mu|$ values, likely making them a good predictive tool. More broadly, it was shown that transition dipoles correlate well with $|\Delta\mu|$ values for all mononitriles and, to a lesser extent, for dinitriles. To a large extent, the correlation for mononitriles is consistent with that expected from the anharmonicity of the nitrile bond. However, Stark effects are larger than just those expected from anharmonicity, suggesting that other contributions are important as well.

The ability to routinely measure vibrational Stark effects to good precision makes it possible to investigate molecular vibrations in condensed phases in a new and sensitive manner. This technique has previously been shown to be useful for measuring an electric field change in a series of myoglobin mutants¹, and has been studied here to clarify the physical origins of vibrational Stark effects.

Acknowledgements

We thank Dr. Arun Chatopadhyay who made early contributions to the VSE of simple nitriles, and Prof. Noel Hush for helpful discussions. This work was supported in part by a grant from the NSF Chemistry Division. The FTIR facilities are supported by the Medical Free Electron Laser Program of the Office of Naval Research under Contract N00014-94-1-1024.

Afterward

Three minor mistakes have been found in this paper since publication, all of which are described in Chapter 7 as well, which has been submitted for publication. *i)* In the discussion of angle dependence this paper says “due to the 3-fold rotational symmetry of acetonitrile, a perpendicular component (of an off-diagonal component) of the transition polarizability cannot occur without significant symmetry breaking by asymmetric solvation.” While it is true that off-diagonal components are forbidden by symmetry, this is not correct for diagonal components, which allows a more satisfactory explanation of the data in Figure 6. *ii)* In the calculation of the anharmonicity contribution to $|\Delta\mu|$, the slopes of the best fit to the data was compared to that expected for anharmonicity. A more accurate comparison would just consider the $|\Delta\mu|$ values. *iii)* In the same calculation, a slope of $0.012 \text{ \AA}/f$ was compared to a slope of 0.0078 \AA , using a f value of 1.1. In this comparison, the Stark effect data was corrected by dividing by f , whereas it should have been multiplied by f .

References and Notes

- (1) Park, E. S.; Andrews, S. S.; Hu, R. B.; Boxer, S. G. *J. Phys. Chem. B* **1999**, *103*, 9813-9817.
- (2) Park, E. S.; Thomas, M. R.; Boxer, S. G. *J. Am. Chem. Soc.* **2000**, *122*, 12297-12303.
- (3) Onsager, L. *J. Am. Chem. Soc.* **1936**, *58*, 1486-1493.
- (4) Stryer, L. *Biochemistry*; Fourth ed.; W. H. Freeman and Company: New York, 1995.
- (5) Laberge, M.; Sharp, K. A.; Vanderkooi, J. M. *J. Phys. Chem. B* **1997**, *101*, 7364-7367.
- (6) Phillips, G. N. J.; Teodoro, M.; Li, T.; Smith, B.; Gilson, M. M.; Olson, J. S. *J. Phys. Chem. B* **1999**, *103*, 8817-8829.
- (7) Kolling, O. W. *J. Phys. Chem.* **1996**, *100*, 16087-16091.
- (8) Nyquist, R. A. *App. Spect.* **1990**, *44*, 1405-1407.
- (9) Spitzer, R. C.; Sievers, A. J.; Silsbee, R. H. *J. Opt. Soc. Am. B* **1992**, *9*, 978-981.
- (10) Lambert, D. K. *J. Chem. Phys.* **1988**, *89*, 3847-3860.
- (11) AM1 level calculations using MacSpartan (Wavefunction Inc.).
- (12) Charles, S. W.; Cullen, F. C.; Owen, N. L. *J. Molec. Struct.* **1976**, *34*, 219-228.
- (13) DeLeon, R. L.; Muenter, J. S. *J. Chem. Phys.* **1984**, *80*, 3992-3999.
- (14) Kumar, A. P.; Rao, G. R. *Spectrochimica Acta Part A* **1997**, *53*, 2033-2039.
- (15) Michel, H.; Lippert, E. *Acetonitrile—The structure of the liquid and solid phases and the nature of the liquid-solid phase transition*; John Wiley & Sons: New York, 1978, pp 293-308.
- (16) Pace, E. L.; Noe, L. J. *J. Chem. Phys.* **1968**, *49*, 5317-5325.
- (17) Reimers, J. R.; Hall, L. E. *J. Am. Chem. Soc.* **1999**, *121*, 3730-3744.

- (18) Thomas, B. H.; Orville-Thomas, W. J. *J. Molec. Struct.* **1971**, 7, 123-135.
- (19) Hush, N. S.; Reimers, J. R. *J. Phys. Chem.* **1995**, 99, 15798-15805.
- (20) Andrews, S. S.; Boxer, S. G. Submitted to *J. Phys. Chem. A* **2001**.
- (21) Mathies, R. A. Experimental and theoretical studies on the excited electronic states of some aromatic hydrocarbons through electric field perturbation and through chemical substituents; Cornell University: Ithaca, NY, 1974.
- (22) Reimers, J. R.; Zeng, J.; Hush, N. S. *J. Phys. Chem.* **1996**, 100, 1498-1504.
- (23) Wortmann, R.; Bishop, D. M. *J. Chem. Phys.* **1998**, 108, 1001-1007.
- (24) Bublitz, G. U.; Boxer, S. G. *Annu. Rev. Phys. Chem.* **1997**, 48, 213-242.
- (25) Jucks, K. W.; Miller, R. E. *J. Chem. Phys.* **1988**, 88, 6059-6067.
- (26) Chattopadhyay, A.; Boxer, S. G. *J. Am. Chem. Soc.* **1995**, 117, 1449-1450.
- (27) Andrews, S. S.; Boxer, S. G. *accepted for Rev. Sci. Inst.* **2000**.
- (28) Long, F.; Freedman, T. B.; Hapanowicz, R.; Nafie, L. A. *Appl. Spectrosc.* **1997**, 51, 504-507.
- (29) Early experiments were performed in a miniature Joule-Thompson refrigerator (MMR Technologies) operating at 80 K, but we found significant sample orientation in an electric field, leading to the development of the immersion cryostat. Poling may not be a problem for AC electric fields.
- (30) Data fitting was done with Macintosh software created for this purpose. The software and C language source code are available upon request.
- (31) The parameters of a modified Voigt are: peak height, frequency, full width at half maximum, and the Gaussian:Lorentzian ratio. These parameters are adjusted only to obtain a best fit to the absorption spectrum, and are not considered to have any physical meaning.
- (32) Assuming the maximum possible θ angle of 90° and the maximum sample rotation angle of $\theta=20^\circ$, C_θ would be corrected to be 13% too small under the

- one-dimensional assumptions, contributing to a 7% error in the calculation of $|\Delta\mu|$.
- (33) Atkins, P. W. *Physical Chemistry*; 5 ed.; W.H. Freeman and Company: New York, 1994.
- (34) Smith, A. L.; Keller, W. E.; Johnston, H. L. *Phys. Rev.* **1950**, *79*, 728.
- (35) Linear coupling between two modes results in redefined normal modes and a separation of the absorption peaks, whereas non-linear coupling gives absorption between the peaks, and is responsible for Fermi resonance. The excess absorption is analogous to the coalescence seen in NMR spectra of dynamic systems (see *e.g.* Gasparro, F.P.; Kolodny, N.H. *J. Chem. Educ.* **1977**, *54*, 258-261).
- (36) Miller, T. M. *Atomic and molecular polarizabilities*; 71 ed.; CRC Press: Boca Raton, FL, 1990, pp 10-193.
- (37) The transition state enthalpy is expected to be about 16 kJ/mol, from butane data (see *e.g.* Kemp, D.S.; Vellaccio, F. *Organic Chemistry*; Worth Publishers: New York, 1980, p. 472), implying an interconversion rate below 1 ms down to the 100K glass transition temperature, which is fast compared to the freezing rate. Thus, low temperature population ratios were computed for 100K, with the assumption that the sample is at equilibrium before freezing and immobilized thereafter. Boltzmann population calculations use an enthalpy difference between *anti* and *gauche* conformations of 2.8 kJ/mol, a value found by AM1 calculations for the relevant nitriles.
- (38) Lowry, T. H.; Richardson, K. S. *Mechanism and Theory in Organic Chemistry*; third ed.; Harper Collins: New York, NY, 1987.
- (39) Cyanogen was synthesized by an adaptation of a method from: Janz, George J., *Inorganic Syntheses*, 5:43-48, 1957.
- (40) McQuarrie, D. A. *Quantum Chemistry*; University Science Books: Sausalito, CA, 1983.
- (41) Strey, G.; Mills, I. M. *Molec. Phys.* **1973**, *26*, 129-138.